

# Prevention Extended Indicator Reference Sheets

## Table of Contents:

TPT_ENROLL: Number of TPT initiations	2
TPT_CON_ENROLL: Number of TPT initiations among contacts	2
TPT_CON04_ENROLL: Number of TPT initiations among contacts <5	2
TPT_PLHIV_ENROLL: Number of TPT initiations among PLHIV	2
PCT_TPT_ADR: Percentage of people with adverse reactions to TPT	2
PCT_TPT_LTFU: Percentage of people started on TPT who were LTFU	3
PCT_TPT_TB: Percentage of people who developed TB while on TPT	3
PCT_TPT_AST: Percentage of people started on TPT with baseline AST/ALT tests	4
TPT_COMPL: Number of people who completed TPT	4
TPT_CON04_COMPL: Number of contacts <5 who completed TPT	4
TPT_CON5PLUS_COMPL: Number of contacts >5 who completed TPT	5

The color of the indicator reference sheet indicates whether the indicator is calculated or not calculated, and whether it could be considered a disaggregate of another indicator:

### Calculated indicators:

Indicator name	<i>Ex: PCT_TB_ART: Percentage TB clients who start or continue ART</i>
----------------	--

### Non calculated indicators:

Indicator name	<i>Ex: TB_ART_N: Number of TB clients who start or continue ART</i>
----------------	---

### Calculated indicators that could be reported as a disaggregate of another indicator:

Indicator name	<i>Ex: PCT_DRTB_ART: Percentage DR-TB clients who start or continue ART</i>
----------------	---

### Non calculated indicators that could be reported as a disaggregate of another indicator:

Indicator name	<i>Ex: DRTB_ART_N: Number of DR-TB clients who start or continue ART</i>
----------------	--

<b>Indicator name</b>	<b>TPT_ENROLL: Number of TPT initiations</b>
<b>Note:</b>	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

<b>Indicator name</b>	<b>TPT_CON_ENROLL: Number of TPT initiations among contacts</b>
<b>Note:</b>	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

<b>Indicator name</b>	<b>TPT_CON04_ENROLL: Number of TPT initiations among contacts &lt;5</b>
<b>Note:</b>	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

<b>Indicator name</b>	<b>TPT_PLHIV_ENROLL: Number of TPT initiations among PLHIV</b>
<b>Note:</b>	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

<b>Indicator name</b>	<b>PCT_TPT_ADR: Percentage of people with adverse reactions to TPT</b>
<b>Note:</b>	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

Indicator name	PCT_TPT_LTFU: Percentage of people started on TPT who were LTFU [Previously PT-13]	
<b>Definition</b>	<p>Percentage of people who started on TB preventive treatment (TPT) who were lost to follow-up (LTFU), among those who started TPT during the reporting period.</p> <p><i>TPT lost to follow-up times are determined by WHO based on the type of TPT regimen. For example, a person is LTFU if TPT interrupted eight consecutive weeks or more for 6H (6 or 9 months of daily isoniazid monotherapy); four consecutive weeks or more for 3HP (3 months of rifapentine plus high dose isoniazid weekly), 3HR (3 months of daily rifampicin plus isoniazid), and 4R (4 months of daily rifampicin); and 10 consecutive days for 1HP (1 month of rifapentine plus isoniazid daily).</i></p>	
<b>Numerator</b>	Number of people who started on TPT and were lost to follow-up (LTFU) during the reporting period.	<i>PBMEF data element: TPT_LTFU</i>
<b>Denominator</b>	Number of people who started TPT during the reporting period.	<i>PBMEF data element: TPT_ALL_ENROLL</i>
<b>Category</b>	Prevent	
<b>Indicator type</b>	Output	
<b>PBMEF level</b>	Extended	
<b>Unit of measure</b>	Percent of people	
<b>Disaggregate by</b>	Risk group (contacts <5 years, contacts ≥5 years, PLHIV), TPT regimen type	

Indicator name	PCT_TPT_TB: Percentage of people who developed TB while on TPT [Previously PT-15]	
<b>Definition</b>	Number of people started on TB preventive treatment (TPT) who developed TB while on TPT during the reporting period	
<b>Numerator</b>	Number of people started on TB preventive treatment (TPT) who developed TB while on TPT during the reporting period	<i>PBMEF data element: TPT_TB</i>
<b>Denominator</b>	Number of people who started TPT during the reporting period.	<i>PBMEF data element: TPT_ALL_ENROLL</i>
<b>Category</b>	Prevent	

<b>Indicator type</b>	Output
<b>PBMEF level</b>	Extended
<b>Unit of measure</b>	Percent of people
<b>Disaggregate by</b>	Risk group (contacts <5 years, contacts ≥5 years, PLHIV), TPT regimen type

<b>Indicator name</b>	<b>PCT_TPT_AST: Percentage of people started on TPT with baseline AST/ALT tests</b> <i>[Previously PT-16]</i>	
<b>Definition</b>	Percentage of people started on TB preventive treatment (TPT) with baseline AST/ALT tests during the reporting period.	
<b>Numerator</b>	Number of people started on TB preventive treatment (TPT) with baseline AST/ALT tests during the reporting period.	<i>PBMEF data element: TPT_AST</i>
<b>Denominator</b>	Number of people who started TPT during the reporting period.	<i>PBMEF data element: TPT_ALL_ENROLL</i>
<b>Category</b>	Prevent	
<b>Indicator type</b>	Output	
<b>PBMEF level</b>	Extended	
<b>Unit of measure</b>	Percent of people	
<b>Disaggregate by</b>	Risk group (contacts <5 years, contacts ≥5 years, PLHIV), TPT regimen type	

<b>Indicator name</b>	<b>TPT_COMPL: Number of people who completed TPT</b>
<b>Note:</b>	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

<b>Indicator name</b>	<b>TPT_CON04_COMPL: Number of contacts &lt;5 who completed TPT</b> <i>[Previously PT-10]</i>
<b>Definition</b>	Number of contacts <5 years who completed TB preventive treatment (TPT) during the reporting period.

<b>Numerator</b>	Number of contacts <5 years who completed TB preventive treatment (TPT) during the reporting period.	<i>PBMEF data element: TPT_CON04_COMPL</i>
<b>Denominator</b>	N/A	<i>N/A</i>
<b>Category</b>	Prevent	
<b>Indicator type</b>	Output	
<b>PBMEF level</b>	Extended	
<b>Unit of measure</b>	Number of children <5	
<b>Disaggregate by</b>	This indicator could be reported as a disaggregate of TPT_COMPL.	

<b>Indicator name</b>	<b>TPT_CON5PLUS_COMPL: Number of contacts &gt;5 who completed TPT</b> <i>[Previously PT-11]</i>	
<b>Definition</b>	Number of contacts aged 5 years and above who completed TB treatment during the reporting period.	
<b>Numerator</b>	Number of contacts aged 5 years and above who completed TB treatment during the reporting period.	<i>PBMEF data element: TPT_CON5PLUS_COMPL</i>
<b>Denominator</b>	N/A	<i>N/A</i>
<b>Category</b>	Prevent	
<b>Indicator type</b>	Output	
<b>PBMEF level</b>	Extended	
<b>Unit of measure</b>	Number of people	
<b>Disaggregate by</b>	This indicator could be reported as a disaggregate of TPT_COMPL.	